

Hypoglycemic Effects in Response to Abelmoshus Esculentus Treatment: A Research Framework using STZ-Induced Diabetic Rats

Indah Mohd Amin

Abstract—*Abelmoshus esculentus*, AE (okra or lady's finger) is flowering plant and cultivated throughout the tropical and temperate region in the world. This plant is popular with its mucilaginous properties and has been acclaimed to have various health benefits which include anti-diabetic properties. This paper will provide the overview of the research framework and giving insight of the experimental procedure to be implemented to investigate the differential expression analysis of liver tissue in streptozotocin-induced diabetic rat in response to *Abelmoshus esculentus* (AE) treatment. The hypoglycemic effects of water extract prepared from the fruit of AE will be studied in diabetic rats (streptozotocin induced). The differential expression study will be carried out by using real time RT-PCR method. Four diabetes-specific genes of interest (carboxylesterase 2, stearoyl-Coenzyme A desaturase 1, insulin-like growth factor 1 and insulin-like growth factor binding protein 2 binding protein) will be selected and the expression level of these genes will be examined quantitatively. The abnormal expression of genes in STZ group will be rescued by the AE therapy. The expected findings from both *in vivo* and molecular studies may reveal the anti-diabetic properties of the AE and suggest that the plant extract may be useful for the management of the disease. A review of relevant literature and studies on AE will also be presented in this paper.

Index Terms— diabetes mellitus, *abelmoshus esculentus*, gene expression, streptozotocin

I. INTRODUCTION

Diabetes can be described as the increase of glucose (sugar) in the blood. It is important to prevent diabetes complications with careful monitoring of blood sugars. Diabetes is correlated with abnormalities in variety of microvascular, macrovascular, neurologic and infectious complications including carbohydrate, fat, and protein metabolism. Type 2 DM begins with a period of insulin resistance with increased pancreatic insulin secretion. As the disease progresses, pancreatic function weaken and as a result, insulin levels fail to keep up with the body requirements [1].

Currently DM is considered one of the chronic diseases and is a condition that is increasing in epidemic proportions throughout the world. WHO reported that diabetes affects about 5% of the global population [2]. The management of diabetes without any side effects is still a challenge to the

medical system as the treatment for diabetes is relatively limited with significant side effects. There is growing interest in the use of natural health products as an alternative approach to current medications. Plant sources has become a target to explore new drugs and in searching biologically active compounds [3].

The streptozotocin-induced diabetic rat is still considered as an important means for the pathophysiology and pharmacology studies of diabetes mellitus. Some of the research has done using diabetic rat to investigate the anti-diabetic properties are mentioned in the literature. The next section discusses the literature review, research framework, expected results and conclusions.

II. LITERATURE REVIEW

This section provides a summary of the literature review on (i) diabetes mellitus, (ii) AE, (iii) scientific study on anti-diabetic properties of medicinal plants from different countries, (iv) research on bioactive properties of AE, (v) chemical and nutritional constituents of *Abelmoshus esculentus*, (vi) nutritional healing properties of AE and (vii) AE-based outcome studies.

A. Diabetes Mellitus (DM)

TABLE 1. COMPARING TYPE 1 AND TYPE 2 DIABETIC MELLITUS [6]

Feature	Type 1 DM	Type 2 DM
Frequency	10-20%	80-90%
Age at onset	Early (below 35 years)	Late (after 40 years)
Type of onset	Abrupt and severe	Gradual and insidious
Weight	Normal	Obese/non-obese
Family history	<20%	About 60%
Genetic locus	Unknown	Chromosome 6
Pathogenesis	Autoimmune destruction of β -Cells	Insulin resistance, impaired insulin secretion
Islet cell antibodies	Yes	No
Blood insulin level	Decreased insulin	Normal or increased insulin
Islet cell changes	Insulinitis, β -cell depletion	No insulinitis, later fibrosis of islets.
Clinical management	Insulin and diet	Diet, exercise, oral drugs, insulin
Acute complications	Ketoacidosis	Hyperosmolar coma

Diabetes mellitus (DM) is the most challenging metabolic disorder as it cannot be cured but needs only to be managed.

Manuscript received April 4, 2011

Indah Mohd Amin is currently with the Faculty of Dentistry, Universiti Teknologi MARA (UiTM) Malaysia, 40450, Shah Alam, Selangor, Malaysia. (Phone: +603-55435836; fax: +603-55435803; e-mail: indahma@salam.uitm.edu.my).

In Malaysia, it is reported in 2010, that the diabetic sufferers are approximately 3.4 million [4]. By far, Asia was recognized as having the potential increase with 2.5 to 3 times more common diabetes [4]. Hence, by 2010, Asia was expected to have 138 million diabetic patients [4].

There are many clinical and pathophysiologic differences between Type 1 and Type 2 diabetes mellitus (DM) and the contrasting features is clearly seen in Table 1 [5, 6].

B. *Abelmoschus Esculentus* (AE)

According to Maganha *et. al* (2009) [7] AE, known as lady’s finger, is a very poorly studied species. It is reported that the biological activity in this species was the *in vitro* antioxidant potential [8]. The major antioxidant molecules were identified to be *quercetin* derivatives and *-epigallocatechin* [8]. There is lacking scientific reported study on these plant properties despite its wide usage as medicinal plant. Therefore, the main objective of this study is to investigate the underlying molecular mechanism on the natural hypoglycemic substances effect in AE.

C. Scientific study on anti-diabetic properties of medicinal plants from different countries

TABLE 2. STUDY ON ANTI-DIABETIC PROPERTIES OF MEDICINAL PLANTS

Plant	Country	Author
<i>Spergularia purpurea</i>	Morocco	[8]
<i>Ferula persica</i> , <i>Paronychia argentea</i> , and <i>Pistacia atlantica</i>	Jordan	[9]
<i>Piper sarmentosum</i>	Thailand	[10, 11]
<i>Pandanus odoratus</i>		
<i>Pseuderanthemum palatiferum</i>	Vietnam & Thailand	[12]
<i>Phellinus baumii</i>	Asean	[13]
<i>Buchholzia coriacea</i>	Southern Nigeria, Africa	[14]
<i>Cymbopogon citratus</i> <i>Stapf.</i>	Nigeria	[15]
<i>Sclerocarya birrea</i>	South Africa	[16]
<i>Musanga cecropioides</i>	Africa	[17]
<i>Cogniauxia podoleana</i>	Congo	[18]

Table 2 provides a summary of the literature on the study relevant to anti-diabetic properties of medicinal plants from various countries.

D. Research on Bioactive Properties Of AE

TABLE 3. RESEARCH ON BIOACTIVE PROPERTIES OF AE

Bioactive Properties	Author
Antitumor cancer	[19]
Antiproliferative & Proapoptotic	[20]
Antioxidant	[21, 22, 23, 24]
Antihemolytic & antioxidant	[25]
Anti-adhesive & antibacteria	[26, 27]
Mucilaginous effect	[28]
Hypocholesterolemic effect	[29]
Inhibitory activities against certain intestinal disaccharidases	[30]
Antiallergic	[31]
Hypoglycemic effect	[32]

Table 3 presents the literature on the research on bioactive properties of AE.

E. Chemical and nutritional constituents of *Abelmoschus esculentus*

TABLE 4. CHEMICAL AND NUTRITIONAL CONSTITUENTS OF AE

Constituents	Author
Uracil, uridine, deoxyuridine, adenine, Adenosine, 3'-deoxyadenosine, Guanosine, deoxyguanosine, phenylalanine Tyrosine, leucine, and isoleucine	[33]
β-carotene	[34]
Protein, fat, fiber, amino acid	[35]
Phospholipids fraction	[36]
Triacylglycerols component	[37]
Ash, carbohydrate, protein, moisture, fat, fiber contents & etc	[38]
Volatile components	[39]

Table 4 presents the literature on chemical and nutritional constituents of AE.

F. Nutritional Healing Properties of AE

TABLE 5. NUTRITIONAL HEALING PROPERTIES OF AE

Healing properties	Author
Anti-ulcer food	[40]
Anticancer, reduced heart attack, lower blood cholesterol, relieve intestinal disorder, relieve inflammation of the colon, relieve diverticulitis, relieve stomach ulcer	[41]
Neutralize acid, lubricate large intestine, treatment of lung inflammation, treatment of irritable bowel, keep joints limber, treatment of sore throats, heal burn soothe poison, soothe psoriasis	[42]

Table 5 presents the literature on nutritional healing properties of AE.

G. AE-Based Outcome Studies

TABLE 6. AE-BASED OUTCOME STUDIES

AE-Based Outcome	Author
Colon targeted tablets	[43]
Oral sustained release matrix tablets	[44, 45]
Disintegrant tablets	[46]
Food additive to produce bread	[47]
Fat replacer in chocolate bar cookies	[48]

Table 6 presents the literature on different results of different studies which are based on AE.

III. RESEARCH FRAMEWORK

The research framework of this study comprise of; preparation of the aqueous extract, animal experiments, isolation of total RNA, real time reverse transcriptase-PCR (RT-PCR), gel electrophoresis.

The following provides the explanation of each of the stages:

A. Preparation Of The Aqueous Extract

The plant in this study will be collected in its natural habitats and botanically determined by Forestry Research Institute Malaysia (FRIM), Malaysia. The whole plant will be ground and 50 g of powder mixed with 500 ml of distilled water (10%). The mixture is heated and boiled under reflux for 30 min. The decoction obtained is centrifuged, filtered, frozen at -20 °C, lyophilised and used without further purification.




Figure 1. Research Framework

B. Animal Experiments

Experiments will be carried out in adult male Sprague-Dawley rats weighing from 130 to 150 g at age of 5 weeks according to standard procedure. Animals house under standard environmental conditions ($23 \pm 1^\circ\text{C}$, $55 \pm 5\%$ humidity and a 12-h light/dark cycle) and maintained with free access to water and a standard laboratory diet. Experimental rats will be induced by intravenous injection of streptozotocin (STZ) (Sigma, St. Louis, MO, USA) into the tail vein (dose 50 mg/kg body weight). STZ is dissolved in 0.1M cold sodium citrate buffer, pH 4.5 and the control rats are injected with vehicle alone. Diabetes will be verified by evaluating blood glucose levels and diabetic rats are confirmed if blood glucose level greater than 300mg/dl (16.7 mmol/l). All the animals are randomly divided into three groups: ($n=7$ per group).

Group 1: Normal healthy control group (N group), normal rats received 0.9% NaCl solution.

Group 2: STZ-induced diabetic group (STZ group), diabetic rats treated with 0.9% NaCl solution

Group 3: AE-treated diabetic group (AE group), diabetic rats treated with AE at three different level (100, 150 and 200mg/kg body weight) and blood glucose levels determine in fasted rats, every 4 to 6 hrs, after 2 and 4 days, 1 and 2 weeks of single or repeated oral administration.

C. Isolation of Total RNA

Total RNA from frozen liver tissue will be isolated with the RNeasy Mini Kit (Qiagen, Stanford, CA). Samples will be processed following the manufacturer's directions. At the last step, the RNA will be eluted with 50 μl of RNase-free

water. Quality of the RNA will be assessed using Agilent Bioanalyzer System.

D. Real Time Reverse Transcriptase-PCR (RT-PCR)

The assay performs using a one-step RT-PCR kit (Qiagen, Germany). A set of primers and a probe will be selected for the in-house real-time RT-PCR assay, forward, reverse, and TaqMan probe. The in-house real-time TaqMan assay will be carried out using QuantiTect Probe RT-PCR kit (Qiagen, Germany). The reaction contained 5 μl of RNA, 0.6 μM of each primer, 0.2 μM of the TaqMan probe. An Applie Biosystem real time-PCR machine is used with the following thermal steps: reverse transcription 30 min, initial denaturation, followed by 40 cycles of denaturation, annealing for , and extension

E. Gel Electrophoresis

Assessment of PCR products will be carried out using Agilent Bionalyzer system. The PCR product will be visualized as a single compact band with expected size.

IV. EXPECTED RESULTS

The following subsections provide insights on the expected results from the experiments.

A. Preparation Of The Water Extract

In this study, a water extract use as a method of choice in order to follow similar traditional preparation had done in many countries. Moreover it also used as a preliminary study before searching on the pure compound.

B. Animal Experiments

A comparison of the glucose levels from three different groups will be studied. STZ induce diabetic rat should significantly elevated the blood glucose levels after 3 months compared with rats injected with vehicle alone (control group). These observations are according to Verspohl (2002) [49], for the development of a type 2 diabetic model. In the diabetic rat model, the extracts will show significant hypoglycemic effect and the effect of the administered extract is dose dependant. It is expected that the lower dose (100 mg/kg) of the AE only produced a statistical significant effect less than the higher dose (150 mg/kg), after its administration.

The data obtained from this study will be the first documented data demonstrated the appropriate doses that can be traditionally used for treating diabetic people. In addition, the result should not show any toxicity and mortality occur to the control group. This is based on the fact that there is no data reported of toxicity occur among its consumer.

C. Molecular Studies

Gene expression can be assessed by measuring the quantity of the final product for instance, the protein. However in this study, measuring RNA level is chosen because it is more efficient with the current technologies and easy access of information. Moreover, available proteomic technologies are generally lower throughput and more challenging.

It is critically important to have a good quality of RNA in

any of expression study. To ensure the reliability of the RNA samples, RNA assessment will be carried out using Agilent Bioanalyzer system. Successful isolation technique yields good quality of RNA. The intact total RNA sample will show distinct 18S and 28S subunit spikes, with 2:1 ratio (28:18S).

In this study, real-time RT-PCR is chosen because it enables us to view PCR amplification in each cycle in faster and easiest way. Compare to other quantitative RT-PCR, real time RT-PCR requires shorter development, normally less variable and more reproducible. Protocol chose for this method is using Taq-Man assay. The advantage of this method is more specific because of the probe designed and will also avoid the primer dimmer.

In studies of target gene expression by RT-PCR, the use of internal reference genes is required to control for RNA quality, reverse transcription efficiency and overall transcriptional activity in samples. Preferably, internal standards should be constitutively expressed by all cell types independent of experimental conditions and they should not be affected by any disease. For normalization, housekeeping genes of β -actin is employed. From this study, the expected real time RT-PCR product will be confirmed by post-PCR gel analysis. This additional step will be carried out as to make sure the amplified product within the expected size

Analysis of gene profiling study on STZ induce diabetic rat did by Hwang et al, 2009 [50], demonstrated many changes occurred in their gene expression. More than a half of gene analyzed (565 from total 835 genes) showing two-fold differences compared to the control. The differentially expressed genes involved in carbohydrate, amino acid metabolism, immunity and defense, lipid, fatty acid, steroid metabolism and signal transduction [50].

In this study, four specific diabetes genes will be chosen to present the hypoglycaemic effect of the extract. Treatment with AE may show significant changes in rescue the abnormal expression of genes in STZ induced diabetic rat.

V. CONCLUSIONS

This paper provides insights of the research framework to investigate the differential expression analysis of liver protein in streptozotocin-induced diabetic rat in response to AE treatment. The expected findings from both *in vivo* and molecular studies may reveal the anti-diabetic properties of the AE and suggest that the extract could be developed as a prospective phytochemical plant. The risks involve in this study may include the animals experimental ranging from ethical consideration, pre, during and after experiments being conducted. The physiological status of each of the rat may give different results with different interpretation.

One limitations of the experiment is on the molecular approach. Using four genes to represent the underlying molecular may not be sufficient. Therefore, future works may involve larger number of genes platform system, for example using microarray as a tool to reveal the molecular mechanism of the hypoglycemic of AE. The findings of the expected results from this experiment may provide indications of the changes in gene expression that may reveal underlying mechanism of diabetic pathophysiology, suggesting new potential target for drug discovery.

ACKNOWLEDGMENT

This research is funded by the Excellence Fund of Research Management Institute, Universiti Teknologi MARA (UiTM) Malaysia. The author would like to thank UiTM Malaysia for its financial and support in completing this work.

REFERENCES

- [1] S. Inzucchi, "Classification and diagnosis of diabetes mellitus," in *Elenberg & Rifkin's Diabetes Mellitus*, D. Porte, R. Sherwin, A. Baron, Eds. New York: McGraw Hill, 2003, pp. 274.
- [2] WHO, *Traditional medicine strategy 2002–2005*, WHO Publications, 2002, pp 1-6.
- [3] G. Y. Yeh, D. M. Eisenberg, D. M. Kaptchuk, and T. J. Phillips "Systematic review of herbs and dietary supplements for glycemic control in diabetes," *Diabetes Care*, vol. 26, 2003, pp. 1277–1294.
- [4] Rugayah, B., The Second National Health and Morbidity Survey II - Diabetes Epidemic in Malaysia was published in 1997 by the Public Health Institute, Ministry of Health, and Malaysia., 1997. <http://www.diabetes.org.my/article.php?aid=63nik>
- [5] E. Brown, *Pathology: PreTest self-assessment and review*, McGraw-Hill Professional, 2004.
- [6] H. Mohan, *Essential Pathology for Dental Students*, Jaypee Brothers Publishers, 2005.
- [7] E. G. Maganha, R. d. C. Halmenschlager, R. M. Rosa, J. A. P. Henriques, A. L. L. d. P. Ramos and J. Saffi, "Pharmacological evidences for the extracts and secondary metabolites from plants of the genus Hibiscus," *Food Chemistry*, vol. 118, no. 1, 2010, pp. 1-10,
- [8] G. Shui and L. L. Peng. An improved method for the analysis of major antioxidants of Hibiscus esculentus Linn. *Journal of chromatography*, 2004, vol. 1048, pp. 17–24.
- [8] M. Eddouks, H. Jouad, M. Maghrani, A. Lemhadri and R. Burcelin, "Inhibition of endogenous glucose production accounts for hypoglycemic effect of Spergularia purpurea in streptozotocin mice," *Phytomedicine*, vol. 10, 2003, no. 6-7, pp. 594-599.
- [9] I. I Hamdan and F. U. Afifi, "Studies on the in vitro and in vivo hypoglycemic activities of some medicinal plants used in treatment of diabetes in Jordanian traditional medicine," *Journal of Ethnopharmacology*, vol. 93, no. 1, 2004, pp. 117-121.
- [10] P. Peungvicha, S. S. Thirawarapan, R. Temsiririrkkul, H. Watanabe, J. K. Prasain and S. Kadota, "Hypoglycemic effect of the water extract of Piper sarmentosum in rats," *Journal of Ethnopharmacology*, vol. 60, no. 1, 1998, pp. 27-32.
- [11] P. Peungvicha, R. Temsiririrkkul, J. K. Prasain, Y. Tezuka, S. Kadota, S. S. Thirawarapan and H. Watanabe, "4-Hydroxybenzoic acid: a hypoglycemic constituent of aqueous extract of Pandanus odoratus root," *Journal of Ethnopharmacology*, vol. 62, no. 1, 1998, pp. 79-84.
- [12] P. Padee, S. Nualkaew, C. Talubmook, S. Sakuljaitrong, "Hypoglycemic effect of a leaf extract of Pseuderanthemum palatiferrum (Nees) Radlk. in normal and streptozotocin-induced diabetic rats," *Journal of Ethnopharmacology*, vol. 132, no. 2, 2010, pp. 491-496.
- [13] H.-J. Hwang, S.-W. Kim, J.-M. Lim, J.-H. Joo, H.-O. Kim, H.-M. Kim and J. W. Yun, "Hypoglycemic effect of crude exopolysaccharides produced by a medicinal mushroom *Phellinus baumii* in streptozotocin-induced diabetic rats," *Life Sciences*, vol. 76, no. 26, 2005, pp. 3069-3080.
- [14] R. A. Adisa, M. I. Choudhary and O. O. Olorunsogo, "Hypoglycemic activity of *Buchholzia coriacea* (Capparaceae) seeds in streptozotocin-induced diabetic rats and mice," *Experimental and Toxicologic Pathology*, 2010.
- [15] A. A. Adeneye and E. O. Agbaje, "Hypoglycemic and hypolipidemic effects of fresh leaf aqueous extract of *Cymbopogon citratus* Stapf. in rats," *Journal of Ethnopharmacology*, vol. 112, no. 3, 2007, pp. 440-444.
- [16] J. A. O. Ojewole, "Hypoglycemic effect of *Sclerocarya birrea* (A. Rich.) Hochst. [Anacardiaceae] stem-bark aqueous extract in rats," *Phytomedicine*, vol. 10, no. 8, 2003, pp. 675-681.
- [17] A. A. Adeneye, O. P. Ajagbonna and O. W. Ayodele, "Hypoglycemic and antidiabetic activities on the stem bark aqueous and ethanol extracts of *Musanga cecropioides* in normal and alloxan-induced diabetic rats," *Fitoterapia*, vol. 78, no. 7-8, 2007, pp. 502-505.
- [18] M. Diatwa, C. B. Samba, T. C. Hondi Assah and A. A. Abena, "Hypoglycemic and antihyperglycemic effects of diethyl ether fraction isolated from the aqueous extract of the leaves of *Cordia alliodora*

- podoleana Baillon in normal and alloxan-induced diabetic rats," *Journal of Ethnopharmacology*, vol. 92, no. 2-3, 2004, pp. 229-232.
- [19] Y. Okada, M. Okada, Y. Sagesaka, "Screening of dried plant seed extracts for adiponectin production activity and tumor necrosis factor-alpha inhibitory activity on 3T3-L1 adipocytes," *Plant Foods For Human Nutrition*, vol. 65, no. 3, 2010, pp. 225-232.
- [20] M. Vayssade, N. Sengkhamparn, R. Verhoef, C. Delaigue, O. Goundiam, P. Vigneron, A. Voragen, H. Schols & M. Nagel, "Antiproliferative and proapoptotic actions of okra pectin on B16F10 melanoma cells," *Phytotherapy Research: PTR*, vol. 24, no. 7, 2010, pp. 982-989.
- [21] S. Atawodi, J. Atawodi, G. Idakwo, B. Pfundstein, R. Haubner, R. Owen, et al., "Polyphenol composition and antioxidant potential of Hibiscus esculentus L. fruit cultivated in Nigeria," *Journal Of Medicinal Food*, vol. 12, no. 6, 2009, pp. 1316-1320.
- [22] N. Ansari, L. Houlihan, B. Hussain, and A. Pieroni, "Antioxidant activity of five vegetables traditionally consumed by South-Asian migrants in Bradford, Yorkshire, UK," *Phytotherapy Research: PTR*, vol. 19, no. 10, 2005, pp. 907-911.
- [23] R. Nagashree, R. Latha, C. Anand, and V. Karthikeyan, "Effect of Hibiscus esculentus fruit on arsenic induced oxidative changes in albino rats," *Journal of Natural Remedies*, vol. 9, no. 2, 2009, pp. 170-176.
- [24] O. E. Adelakun, O. J. Oyelade, B. I. O. Ade-Omowaye, I. A. Adeyemi, and M. Van de Venter, "Chemical composition and the antioxidative properties of Nigerian Okra Seed (*Abelmoschus esculentus* Moench)," *Flour Food and Chemical Toxicology*, vol. 47, no. 6, 2009, pp. 1123-1126.
- [25] M. A. Ebrahimzadeh, S. F. Nabavi, and S. M. Nabavi, "Antihemolytic and antioxidant activity of Hibiscus esculentus leaves," *Pharmacologyonline*, vol. 2, 2009, pp. 1097-1105.
- [26] N. Wittschier, Lengsfeld C, Vorthems S, Stratmann U, Ernst J, Hensel A, et al., "Large molecules as anti-adhesive compounds against pathogens," *The Journal Of Pharmacy And Pharmacology*, vol. 59, no. 6, 2007, pp. 777-786.
- [27] C. Lengsfeld, F. Titgemeyer, G. Faller, and A. Hensel, "Glycosylated compounds from okra inhibit adhesion of *Helicobacter pylori* to human gastric mucosa," *Journal Of Agricultural And Food Chemistry*, vol. 52, no. 6, 2004, pp. 1495-1503.
- [28] K. Ameena, C. Dilip, R. Saraswathi, P. N. Krishnan, C. Sankar, S. P. Simi, "Isolation of the mucilages from Hibiscus rosasinensis linn. and Okra (*Abelmoschus esculentus* linn.) and studies of the binding effects of the mucilages," *Asian Pacific Journal of Tropical Medicine*, vol. 3, no. 7, 2010, pp. 539-543.
- [29] P. Srinivasa Rao, P. Udayasekhara Rao and B. Sesikeran, "Serum cholesterol, triglycerides and total lipid fatty acids of rats in response to okra (*Hibiscus esculentus*) seed oil," *Journal of the American Oil Chemists' Society*, vol. 68, no. 6, 1991, pp. 433-435.
- [30] H. M. M., Hassan, "Inhibitory activities of some mucilages and gums against certain intestinal disaccharidases," *Australian Journal of Basic and Applied Sciences*, vol. 3, no. 3, 2009, pp. 2741-2746.
- [31] S. Tewtrakul, A. Itharat, P. Thammaratwasik, B. Ooraikul, "Anti-allergic and anti-microbial activities of some Thai crops Songklanakarinn," *Journal of Science and Technology*, vol. 30, no. 4, 2008, pp. 467-473.
- [32] M. Aslan, E. Sezik. and E. Yeşilada, "Effect of hibiscus esculentus L. Seeds on blood glucose levels in normoglycaemic, glucose-hyperglycaemic and streptozotocin-induced diabetic rats Gazi Üniversitesi Eczacılık Fakültesi Dergisi", vol. 20, no. 1, 2003, pp. 1-7.
- [33] L. Jia, H-F., Li, L.-L. Jing, "Chemical constituents in n-butanol extract of *Abelmoschus esculentus* Chinese Traditional and Herbal Drugs", vol. 41, no. 11, 2010, pp. 1771-1773.
- [34] J. M. El-Qudah, "Identification and quantification of major carotenoids in some vegetables," *American Journal of Applied Sciences*, vol. 6, no. 3, 2009, pp. 492-497.
- [35] P. Udayasekhara Rao, "Chemical composition and biological evaluation of Okra (*Hibiscus esculentus*) seeds and their kernels," *Plant Foods for Human Nutrition (Formerly Qualitas Plantarum)*, vol. 35, no. 4, 1985, pp. 389-396.
- [36] S. Fiad, "Phospholipids of six seed oils of malvaceae," *Journal of the American Oil Chemists' Society*, vol. 68, no. 1, 1991, pp. 26-28.
- [37] S. Fiad, "Component triacylglycerols of six seed oils of malvaceae," *Journal of the American Oil Chemists' Society*, vol. 68, no. 1, 1991, pp. 23-25.
- [38] J. Hussain, A. L. Khan, N. U. Rehman, M. Hamayun, T. Shah, M. Nisar, T. Bano, and I.-J. Lee, "Proximate and nutrient analysis of selected vegetable species: A case study of Karak region," *Pakistan African Journal of Biotechnology*, vol. 8, no. 12, 2009, pp. 2725-2729.
- [39] M. Camciuc, J. M. Bessièrè., G. Vilarem, A. Gaset, "Volatile components in okra seed coat," *Phytochemistry*, vol. 48, no. 2, 1998, pp. 311-315.
- [40] H.K. Bakhru, *Healing Through Natural Foods*, 2000, Jaico Publishing House.
- [41] P. A. Balch, *Prescription for Dietary Wellness*, 2003, Avery.
- [42] B. Mars, *Rawsome!: Maximizing Health, Energy, and Culinary Delight With the Raw Foods Diet*, 2004, Basic Health Publications, Inc.
- [43] Ilango, K.B.a, Mishra, M.b, Devi, S.c, Rajsekaran, A.c, Senthil kumar, M.c, Subburaju, T.a In vitro and in vivo evaluation of okra polysaccharide-based colon-targeted drug delivery systems International Journal of Pharmaceutical Sciences Review and Research Volume 5, Issue 1, November 2010, Pages 138-145
- [44] M. Emeje, O. Olaleye, C. Isimi, J. Fortunak, S. Byrn, O. Kunle, and S. Ofoefule, "Oral sustained release tablets of zidovudine using binary blends of natural and synthetic polymers," *Biological and Pharmaceutical Bulletin*, vol. 33, no. 9, 2010, pp. 1561-1567.
- [45] V. D. Kalu, M. A. Odeniyi and K. T. Jaiyeoba, "Matrix properties of a new plant gum in controlled drug delivery," *Archives of Pharmacal Research*, vol. 30, no. 7, 2007, pp. 884-889.
- [46] Bakre, L.G., Jaiyeoba, K.T. Evaluation of a new tablet disintegrant from dried pods of *Abelmoschus esculentus* L (Okra) Asian Journal of Pharmaceutical and Clinical Research Volume 2, Issue 3, July 2009, Pages 83-91
- [47] Acquistucci, R, & Francisci, R 2002, 'Effect of okra (*Hibiscus esculentus* L.) addition on the technological properties of a wheat flour', International Journal Of Food Sciences And Nutrition, 53, 5, pp. 375-379, MEDLINE with Full Text, EBSCOhost, viewed 2 April 2011.
- [48] Romanchik-Cerpovicz, J, Tilmon, R, & Baldree, K 2002, 'Moisture retention and consumer acceptability of chocolate bar cookies prepared with okra gum as a fat ingredient substitute', Journal Of The American Dietetic Association, 102, 9, pp. 1301-1303, MEDLINE with Full Text, EBSCOhost, viewed 2 April 2011.
- [49] Verspohl, E.J., Recommended testing in diabetes research. *Planta Medica*, vol. 68, 2002, pp. 581-590.
- [50] H. J. Hwang, S. W. Kim, Y. M. Baek, S. H. Lee, H. S. Hwang and J. W. Yun, Gene Expression profiling in streptozotocin-induced diabetic rat liver in response to fungal polysaccharide treatment, *Korean J. Chem. Eng.*, vol. 26, no 1, 2009, pp. 115-126.



Indah Mohd Amin was born at Hospital Duchess of Kent, Sandakan, Sabah, Malaysia, on the 9th June 1978. She obtained a degree in molecular and cell biology (biochemistry) from Universiti Kebangsaan Malaysia (UKM), Bangi, Selangor, Malaysia in 2000. Then, she obtained master in science (molecular and medicine) from Universiti Teknologi MARA (UiTM) Malaysia, Shah Alam, Selangor, Malaysia in 2007.

Previously, she was a lecturer at the Faculty of Health and Life Sciences, Management Science University (MSU), Shah Alam, Selangor, Malaysia from the year 2005 to 2008. Currently, she is a biochemistry lecturer at the Faculty of Dentistry, Universiti Teknologi MARA (UiTM) Malaysia, Shah Alam, Selangor, Malaysia, since 2008. Her research interests are in molecular medicine, biochemistry, biotechnology, bioinformatics and health informatics.

Ms Indah Mohd Amin is a member of Institute of Electrical and Electronics Engineers (IEEE), International Association of Computer Science and Information Technology (IACSIT), Asia-Pacific Chemical, Biological and Environmental Engineering Society (APCBES) and International and American Association for Dental Research (IADR).