

Effects of Supplemental Dietary Phytase on Performance and Blood Biochemical Parameter of Broiler Chicks

M. A. Jalali and M. Babaei

Abstract—An experiment was conducted to study the effects of phytase enzyme on performance, blood biochemical parameter and tibia ash content of broiler chicks. Two hundred eighty Ross broiler chicks were located randomly to seven experimental diets at four replicates (cages) and fed from 5 to 28 days of age. A positive control diet (0.422% non-Phytate Phosphorus, nPP) and a negative control diet (0.316% nPP) were used, and five more diets were manufactured by supplementing the negative control diet with 300, 600, 1,200, 2,400, and 24,000 FTU/kg of exogenous phytase (Phyzyme XP). The addition of exogenous phytase above 600 FTU/kg and 1200 FTU/kg improved ($P<0.05$) feed conversion ratio and body weight gain of the birds fed the negative control diet, respectively. Decreasing nPP content in the diet reduced tibia ash ($P<0.001$) content and phytase supplementation increased it ($P<0.001$) as logarithmic liner equation. Phosphorus concentration of serum chicks fed negative control diet was reduced and phytase supplementation above 600 FTU/kg increased it. However, negative control diet increased alkaline phosphatase activity of serum and addition of phytase above 2400 FTU/kg reduced its activity to level of positive control diet ($P<0.01$). Albumin concentration of serum chicks was increased by addition of phytase above 600 FTU/kg diet. The lowest calcium content and highest total protein concentration of serum chicks were seen at treatment of 600 FTU/kg phytase. These results demonstrated that addition of 600 FTU/kg phytase to low-available phosphorus diets (75% of NRC recommended) improved performance of broiler chicks due to improve utilization of phytate phosphorus of corn-soy bean meal diets.

Index Terms—Chicks, phytase, performance, serum metabolites.

I. INTRODUCTION

Phosphorus (P) is a limiting nutrient for algal growth and excess it causes contaminate water by stimulates algal blooms growth and consequently, depletes dissolved oxygen in surface water [1]. Recently the most effective factor in environment contamination is especially contamination by phosphorus [2]. Poultry feeds typically contain a high proportion of cereals, grain legumes, and oilseed meals, which approximately two-thirds of their P content as phytate form [3]. Phytate is an anionic acid that binds P and other cationic substances, which does not allow them to be completely used by bird. Broilers have insufficient enzyme to effectively hydrolyze the phytate P of corn and soybean meal. To meet dietary P requirement of broiler, diets are

supplemented with inorganic P. It and phytate P of broiler diets can be contributed to environmental pollution if the availability of P was low. To help increase the availability of phytate P, exogenous phytase enzymes are being added to commercial diets [4]. Phytase research that has focused on the effect of varying levels of supplemental phytase has shown increases in broiler performance, bone ash, and phytate P utilization with each additional unit or level of phytase [5]. Although many studies have reported improvements in performance parameters associated with the addition of phytases to the diet, few studies have added exogenous phytase at large range of concentrations phytase units (FTU)/kg of diet. On the other hand, few attentions have been done by researcher on effects of phytase supplementation on blood biochemical parameter of chicks. The use of phytase at wide range in broiler diets may be affected health and performance of chicks. Therefore, the aim of this experiment was to evaluate the influence of different levels of dietary supplementation phytase on performance and blood biochemical parameter of broiler chicks.

II. METHODS AND MATERIALS

A. Diets

The National Research Council's (NRC) nutrient values [6] for ingredients were used to formulate the basal diet. The basal corn-soybean meal diet used contained a calculated metabolizable energy of 3000 kcal/g diet (TABLE II) and a deficient non-phytate phosphorus (nPP) level of 0.316% (75% of NRC recommendation) as a negative control diet (C-). To the negative control diet, phytase (Phyzyme XP, Danisco Animal Nutrition, Carol Stream, IL) was added at 300, 600, 1200, 2400, and 24000 FTU/kg. In addition, dicalcium phosphate was added to the basal diet to create a positive control diet (C+) that contained 0.422% nPP (equal NRC recommendation) and no phytase.

B. Birds and Data Collection

Two hundred eighty one-day-old Ross broiler chicks were used for this experiment. Each 10 birds were housed randomly in cages with 1m x 1.2m floor space. Each experimental diet was tested with four replicate cages of ten chicks. Birds from 1 to 5d of age were fed by C+ diet and from 5 to 21 d of age were fed by seven experimental diets. Feed consumption and body weight gain of chicks were recorded and feed conversion ratio (FCR) calculated from 5 to 21d of age. At the end of experiment (21d of age), two chicks were selected randomly from each cage and blood samples were collected from wing vein. Sera were separated and then stored at -20°C until assayed. The right tibia was

Manuscript received January 2, 2012.

The authors are with the Department of Animal Science, Faculty of Agriculture and Natural Resources, Shahrekord Branch, Islamic Azad University, Iran (Tel.: +989132666775, fax: +983813361093, e-mail: Fish.nutritionist@gmail.com).

removed, and ash content of dried, fat free bone was determined as described by AOAC [7]. Concentration of albumin, total protein, calcium, phosphorus and alkaline phosphatase activity of serum were measured by commercial kit based on ZiestChem Diagnostics kits (Tehran, Iran; by Technicon RA1000).

C. Statistical Analysis

All Data were analyzed using the GLM procedure of SAS software [8] for analysis of variance as completely randomized design. Significant differences among treatments were identified at 5% level by Duncan's multiple range tests [9].

TABLE I: COMPOSITION AND CALCULATED ANALYSIS OF NEGATIVE (C-) AND POSITIVE (C+) CONTROL DIET.

Ingredient (%)	negative (C-)	positive (C+)
Ground corn	55.8	55.8
Soybean meal	35.3	35.3
Fish meal	2	2
Sunflower oil	3.201	3.201
Limestone	1.203	1.203
Dicalcium phosphate	0.636	1.2
DL methionine	0.15	0.15
Sand	1.11	0.546
Salt	0.1	0.1
Vitamin mix*	0.25	0.25
Mineral mix*	0.25	0.25
Total	100	100
Calculated analysis		
Metabolizable energy(Kcal/kg)	3000	3000
Crude protein (%)	21.56	21.56
Calcium (%)	0.813	0.937
nPP **(%)	0.316	0.422
Sulphur amino acids (%)	0.844	0.844

* Supplied per kilogram of diet: 18000 IU vitamin A; 4000 IU vitamin D3; 36 IU vitamin E; 4 mg vitamin K3; 3.6 mg vitamin B1; 13.2 mg vitamin B2; 20 mg D-calcium pantothenate; 60 mg niacin; 6 mg vitamin B6; 2 mg folic acid; 30 µg vitamin B12; 0.2 mg biotin; 1000 mg choline chloride; 200 mg Mn; 100 mg Fe; 200 mg Zn; 20 mg Cu; 2 mg I; 0.4 mg Se.

** non-phytate phosphorus

III. RESULTS AND DISCUSSION

The addition of exogenous phytase above 1200 FTU/kg and 600 FTU/kg improved ($P<0.05$) body weight gain (fig. 1) and feed conversion ratio (fig. 2) of the birds fed the negative control diet (C-), respectively. These results are in agreement with other reports [10]. The improve of body weight gain and feed conversion ratio of chicks as a result of phytase application might be due to the increased availability of energy [5] and amino acid digestibility [11] due to a raise in phytate degradation. The ability of phytase to improve P availability by hydrolyzing phytate-bound P in poultry diets

can therefore reduce supplementation of diets with inorganic P sources.

Decreasing nPP content in the diet (negative control, C-) reduced tibia ash ($P<0.001$) content and phytase supplementation increased it ($P<0.001$) as logarithmic liner equation [$Y=2.308 \log (X+I) +44.77$; Y: percent of tibia ash, X: phytase activity of diet (FTU/kg), $R^2=0.74$, $P<0.0001$] (fig. 3). This result is in close agreement with other equation proposed by researcher [5]. Changes in the amount of P available to the animal may affect its bone mineral status. This is the reason for the use of percentage bone ash as an indicator of mineral adequacy in animals [12]. As other experiment carried out by japanese quail, addition of phytase enzyme to the diet improved ash and P of bird bon [13].

TABLE II showed the effects of dietary treatment and graded level of phytase supplement on some blood biochemical parameter. Addition of Phytase enzyme to broiler diet changed blood metabolites. Phosphorus concentration of serum chicks fed negative control diet was reduced and phytase supplementation above 600 FTU/kg increased it. However, negative control diet increased alkaline phosphatase activity of serum and addition of phytase above 2400 FTU/kg reduced its activity to level of positive control diet ($P<0.01$). Other research also showed that the level of alkaline phosphatase in sera chick reduced by phytase [14]. The decrease alkaline phosphatase activity and increased P in serum associated with the diets supplemented with phytase might be reflected the increased availability of phosphorus [15]. It is reported that reduction of phosphorus level in diets, increases alkaline phosphatase activity of broiler chicks [16].

Albumin concentration of serum chicks was increased by addition of phytase above 600 FTU/kg diet (TABLE II). The highest level of albumin and total protein of sera chick were seen at treatment of 1200 FTU/kg phytase. Increasing of albumin and total protein may be related to improve of amino acid availability by phytase [11].

The main objective for supplementing phytase to poultry diets is to improve phytate P use in the birds and reduce the amount of P flowing into the environment in animal waste. These results demonstrated that addition of 600 FTU/kg phytase to low-available phosphorus diets (75% of NRC recommended) improved performance of broiler chicks due to improve utilization of phytate phosphorus of corn-soy bean meal diets. The addition of phytase to broiler diets at levels of greater than 600 FTU/kg diet (until 24000 FTU/kg diet), doesn't have any adverse effects on performance and health of chicks and improved them.

ACKNOWLEDGMENTS

The author is indebted to Dr M. Jafarian for blood analyses, Dr F. Khayrie and S. Tavakoli for assistance in data acquisition.




Fig. 1. Effects of different level of phytase on body weight gain of broiler chicks.

(C+: Positive control diet, C- : Negative control diet.

a-d column values with same superscript are not significantly different (P<0.05).




Fig. 2. Effects of different level of phytase on feed conversion ratio (FCR) of broiler chicks.

(C+: Positive control diet, C- : Negative control diet.

a-d column values with same superscript are not significantly different (P<0.05).




Fig. 3. Effects of different level of phytase on Tibia ash content of broiler chicks.

TABLE II: EFFECTS OF SUPPLEMENTAL DIETARY PHYTASE ON BLOOD BIOCHEMICAL PARAMETER OF BROILER CHICKS.

Phytase level (FTU/kg diet)	Phosphorus (mg/ dl)	Calcium (mg/ dl)	Alkaline phosphatase (U/L)	Albumin (g/ dl)	Total protein (g/ dl)
C+	6.93 ^a	8.96 ^a	407.3 ^b	2.36 ^{bc}	3.63 ^{cd}
C- (0)	4.00 ^c	9.93 ^a	482.3 ^a	1.96 ^c	3.40 ^d
300	3.96 ^c	8.96 ^a	475.0 ^a	1.76 ^c	3.33 ^d
600	6.76 ^{ab}	6.06 ^b	488.0 ^a	3.03 ^{ab}	5.10 ^{ab}
1200	5.16 ^{bc}	7.73 ^{ab}	488.6 ^a	3.46 ^a	5.37 ^a
2400	5.60 ^{abc}	9.13 ^a	420.3 ^b	3.30 ^{ab}	4.16 ^{bcd}
24000	6.30 ^{ab}	8.03 ^{ab}	390.6 ^b	3.26 ^{ab}	4.46 ^{abc}
SEM	0.515	0.709	10.18	0.298	0.294
Probability	0.003	0.035	0.0001	0.0044	0.0009

C+: Positive control diet, C- : Negative control diet.

a-d column values with same superscript or no superscript are not significantly different (P<0.05).

SEM: standard error of mean.

REFERENCES

- [1] Y. Sharple, A. N. T. Ce. Daniel, and D. R. Edwards. "Phosphorus movement in the landscape," *J. Prod. Agric.* 1993, Vol. 6, No. 4, pp. 492–500.
- [2] A. Musapuor, M. Afsharmanesh, and H. Moradi Shahrabak, "Use of microbial phytase for decrease of pollutant due to environmental poultry excreta phosphorus," *Inter. J. of Agri. & Biology.* 2006, Vol. 8, No. 1, pp. 35-37.
- [3] A. J. Cowieson, T. Acamovic, and M. R. Bedford, "Supplementation of corn–soy-based diets with an *Escherichia coli*-derived phytase: effects on broiler chick performance and the digestibility of amino acids and metabolizability of minerals and energy," *Poult. Sci.* 2006, Vol. 85, No. 8, pp. 1389–1397.
- [4] R. L. Payne, T. K. Lavergne, and L. L. Southern, "A comparison of two sources of phytase in liquid and dry forms in broilers," *Poul. Sci.* 2005. Vol. 84, No. 2, pp. 265–272.
- [5] R. B. Shirley and H. M. Jr. Edwards, "Graded levels of phytase past industry standards improves broiler performance," *Poul. Sci.* 2003. Vol. 82, No. 4, pp. 671–680.
- [6] National Research Council, "Nutrient Requirement of Poultry," 9th rev. ed. National Academy Press, Washington, DC. 1994.
- [7] AOAC, Official Methods of Analysis. 15th ed. "Association of Official Analytical Chemists," Arlington, VA. 1990.
- [8] SAS, Institute, Inc., "SAS User's guide," version 6.03 ed., SAS Institute Inc., Cary, North Carolina. 1993.
- [9] D. B. Duncan, "Multiple range and F-Tests," *Biometrics.* 1955. Vol. 11, pp.1-42.
- [10] B. C. Watson, J. O. Matthews, L. L. Southern, and J. L. Shelton, "The effects of phytase on growth performance and intestinal transit time of broilers fed nutritionally adequate diets and diets deficient in calcium and phosphorus," *Poul. Sci.* 2006, Vol. 85, No. 3, pp. 493–497.
- [11] P. H. Selle, V. Ravindran, W. L. Bryden, and T. Scott. "Influence of dietary phytate and exogenous phytase on amino acid digestibility in poultry, areview," *The J. Poul. Sci.*, 2006, Vol. 43, No. 2, pp. 89-103.
- [12] E. M. Onyango, M. R. Bedford, and O. Adeola. "Efficacy of an evolved *Escherichia coli* phytase in diets of broiler chicks," *Poul. Sci.* 2005 Vol. 84, No. 2, pp. 248–255.
- [13] N. Vali and M. A. Jalali, "Influence of different levels of phytase enzyme on japanese quail (*coturnix japonica*) performance," *Agriculturae Conspectus Scientificus.* 2011. Vol. 76, No. 4, pp. 387-390.
- [14] R. Nourmohammadi, S. M. Hosseini, and H. Farhangfar "Effect of citric acid and microbial phytase on serum enzyme activities and plasma minerals retention in broiler chicks," *Afr. J. Biotechnol.* 2011. Vol. 10, No. 62, pp. 13640-13650.
- [15] A. Viveros, A. Brenes, I. Arija, and C. Centeno. "Effects of microbial phytase supplementation on mineral utilization and serum enzyme activities in broiler chicks fed different levels of phosphorus," *Poul. Sci.* 2002. Vol. 81, No. 8, pp. 1172–1183.
- [16] A. Brenes, A. Viveros, I. Arija, C. Centeno, M. Pizarro, and C. Braro, "The effect of citric acid and microbial phytase on mineral utilization in broiler chicks," *Anim. Feed Sci. Technol.* 2003. Vol. 110, No. 1-4, pp. 201-219.